



# Home-garden connectivity rather than tree-cover connectivity facilitates biodiversity in fragmented tropical forest landscapes

Ping Sun · Matthias Baumann · Beth A. Kaplin · Tobias Kuemmerle · Joern Fischer

Received: 21 August 2025 / Accepted: 5 April 2026  
© The Author(s) 2026

## Abstract

**Context** Forest landscape restoration (FLR) has emerged as a global strategy to address forest fragmentation and degradation, yet, its ecological effectiveness at local scales remains uncertain. Further, many FLR interventions focus on tree planting, while overlooking other tree-rich systems such as home gardens.

**Objectives** This study focused on the Afromontane rainforest landscape of western Rwanda—a biodiversity-rich region under intense anthropogenic

pressure. As tree plantations and home gardens represent the dominant land use types in the study area, our research aimed to assess their respective roles in supporting landscape connectivity and biodiversity.

**Methods** First, we evaluated how tree cover and home gardens contribute to landscape connectivity by applying circuit theory through Omniscape. Subsequently, we examined the relationships between tree-cover connectivity, home-garden connectivity, and the richness and diversity of woody plants and birds, using data collected from 91 field sites.

**Results** Tree-cover connectivity was negatively correlated with the richness and diversity of woody plants and birds. This suggested that increased connectivity through mostly exotic trees did not translate into habitat connectivity for biodiversity. In contrast, home-garden connectivity was positively correlated with the richness and diversity of woody plants and birds. This highlighted the possible beneficial role of home gardens as ecological stepping stones.

**Conclusions** We recommend that restoration efforts prioritize the protection of remaining natural forests, which are irreplaceable for sustaining native biodiversity. Complementary FLR strategies should promote the regeneration of secondary forests dominated by indigenous tree taxa and the expansion of biodiverse agroforestry systems, such as home gardens.

---

**Supplementary Information** The online version contains supplementary material available at <https://doi.org/10.1007/s10980-026-02352-9>.

---

P. Sun (✉) · J. Fischer  
Social-Ecological Systems Institute, Faculty  
of Sustainability, Leuphana University Lüneburg,  
Lüneburg, Germany  
e-mail: ping.sun@leuphana.de

P. Sun · M. Baumann · T. Kuemmerle  
Geography Department, Humboldt Universität Zu Berlin,  
Berlin, Germany

P. Sun · B. A. Kaplin  
Center of Excellence in Biodiversity and Natural Resource  
Management, University of Rwanda, KN 7 Ave, Kigali,  
Rwanda

T. Kuemmerle  
Integrative Research Institute on Transformations  
of Human-Environment Systems (IRI THESys),  
Humboldt-Universität Zu Berlin, Berlin, Germany

**Keywords** Biodiversity · Forest landscape restoration · Home gardens · Landscape connectivity · Stepping stones

## Introduction

The growing human population and its demand for agricultural land, mineral resources, and spaces to live have dramatically transformed landscapes (Ellis et al. 2021) and disrupted the ecological networks that support the Earth's life-sustaining ecosystems (Hagen et al. 2012; Rudnick et al. 2012; Haddad et al. 2015; Lewis et al. 2015). Landscape fragmentation, degradation, and unsustainable land use have triggered a range of negative ecological consequences, including declines in habitat connectivity and quality (Noss 1991; Perino et al. 2019), detrimental effects on biodiversity (Sala et al. 2000; Fahrig 2003; Fischer and Lindenmayer 2007; Haddad et al. 2015), and disruptions to both ecosystem functioning (Gonzalez et al. 2009; Peh et al. 2014; Hatfield et al. 2018) and the provision of ecosystem services (Tscharntke et al. 2005; Dobrovolski et al. 2011; Mitchell et al. 2015). These negative impacts are particularly apparent in tropical forest landscapes, which are home to more than half of the Earth's terrestrial biodiversity (Jenkins et al. 2013; Lewis et al. 2015; Gibson et al. 2021). Today, less than half of the native tropical forests remain (Lewis et al. 2015), and once continuous ecosystems have been fragmented into more than 50 million forest patches, producing nearly 50 million kilometers of additional forest edges (Brinck et al. 2017). Most tropical forest fragments are embedded in a complex and dynamic matrix of land uses (hereafter referred to as the 'landscape matrix'; Gascon et al. 1999; Jules and Shahani 2003; Chazdon et al. 2009), where fragments of natural forest are juxtaposed with croplands, pastures, tree plantations, home gardens, and other land uses. Depending on the type of land use, and associated land cover, the landscape matrix can either enhance landscape connectivity and increase biodiversity or, conversely hinder species' movement and lower biodiversity (Ricketts 2001; Kupfer et al. 2006; Ruffell et al. 2017; Boesing et al. 2018; de Souza Leite et al. 2022). Importantly, a 'soft matrix'—with similar vegetation structure to native areas—typically supports ecological processes such as movement and

thus increases overall biodiversity (Anderson et al. 2007; Fischer and Lindenmayer 2007; Chazdon et al. 2009), whereas a 'hard matrix'—with vegetation that greatly differs from native areas—can hinder the movement of many species and causes negative edge effects (Lindenmayer and Fischer 2006).

In recent decades, forest landscape restoration (FLR) has gained traction (Clewell et al. 2004; Maginnis et al. 2012; Chazdon and Guariguata 2018) with the overall aim to foster multifunctional landscapes, balancing ecological integrity and human needs (Lamb 2014; Suding et al. 2015; Reed et al. 2016; Mansourian et al. 2017; Chazdon et al. 2020). In practice, FLR is a broader concept than the traditional notion of 'ecological restoration' (Nyiramvuyekure et al. in review), and it can include many types of ecological interventions, including tree plantations, agroforestry sites, and restoration with native species. Through a combination of interventions, FLR aims to increase tree cover, improve habitat connectivity and biodiversity conservation, enhance landscape resilience to climate change, and improve the well-being of local communities (Brancalion and Chazdon 2017; FAO 2017; Chazdon and Guariguata 2018). Its effectiveness depends on the local context of each landscape (Chazdon et al. 2017; Stanturf et al. 2019), with the spatial configuration of forest fragments and the specific interventions within the landscape matrix being critical parameters for defining and achieving FLR objectives. In the past, FLR interventions in tropical forest landscapes have often involved tree plantations predominantly consisting of non-indigenous tree species. Although tree plantations can support the restoration of basic ecosystem functions (e.g., erosion control, biomass production), they may fall short of effectively protecting biodiversity (Bullock et al. 2011). This is especially the case for commercial tree plantations dominated by monocultures, which are typically established for economic reasons, rather than providing ecological functions and value for biodiversity, especially if they consist of non-native trees (Sayer et al. 2004; Brancalion and Chazdon 2017; Mansourian and Berrahmouni 2021).

In contrast to exotic monoculture plantations, home gardens that incorporate native tree species and culturally important crops can serve as multifunctional landscapes known to increase biodiversity (Sunwar et al. 2006; Engelen et al. 2017; Gbedomon et al. 2017; Panyadee et al. 2018; Rooduijn et al.

2018; Yinebeb et al. 2022), to support local livelihoods (Maroyi 2009; Galhena et al. 2013; Gifawesen et al. 2020; Hansen et al. 2022), and to contribute to both ecological and social resilience (Kumar and Nair 2006). Known as ‘tree home gardens’, traditional home gardens are among the oldest agroforestry systems in tropical and sub-tropical regions (Soemarwoto 1987; Kumar and Nair 2006; Nair et al. 2021). These small but highly diversified micro-ecosystems (Galluzzi et al. 2010) are typically managed by local communities across generations, can harbour rich genetic resources, including some indigenous species that may have gone extinct in degraded natural forests (Montagnini 2020). Under some circumstances, home gardens can also provide foraging habitat for forest specialists (Anderson et al. 2007), contribute to biodiversity conservation (Abebe et al. 2013; Engelen et al. 2017; Rooduijn et al. 2018; Vargas-Cárdenas et al. 2024) and function as ecological corridors or stepping stones (Bhagwat et al. 2008; Campera et al. 2021). How home gardens can contribute to overall landscape connectivity, however, remains unclear.

The Afromontane rainforest landscape of western Rwanda is a tropical biodiversity hotspot with exceptionally high endemism (Plumptre et al. 2007; Mittermeier et al. 2011; Kindt 2014). The region also has the highest human population density in mainland Africa (693 inhabitants per square kilometre; National Institute of Statistics of Rwanda 2023), posing massive pressure on the natural landscape. Historically, the landscape experienced severe deforestation and forest fragmentation, but has also seen extensive restoration efforts over the past decade (World Bank 2014). Rwanda’s FLR initiative aims to protect and restore natural forests, improve access to clean water, enhance woodlot management, and promote agroforestry to realize a multifunctional landscape (Ministry of Natural Resources 2014). According to the national Restoration Opportunities Assessment Methodology (Ministry of Natural Resources 2014), approximately 2.25 million hectares (Mha) of land are suitable for FLR interventions, including 1.5 Mha prioritized for agroforestry, improved silviculture and the establishment of protective forests on steep slopes (erosion-control plantations). To date, large-scale restoration efforts have primarily focused on increasing tree cover, primarily through the establishment of tree plantations in the matrix adjacent to protected areas


(Republic of Rwanda 2012). Almost 90 per cent of restored tree plantations consist of exotic species, and the number of newly planted exotics exceeds the number of native trees manifold (Mugabowindekwe et al. 2023). Moreover, tree plantations are often fragmented, poorly managed and mostly disconnected from natural forests (Nduwamungu 2011). Tree plantations may support some natural biodiversity, however, their role in maintaining biodiversity at the landscape level remains little understood and abides a critical question for restoration and conservation efforts in the region. Conversely, little attention has been allocated to home gardens, despite their potentially high benefits to improve habitat condition, connectivity and ecosystem services provisioning.

Against this background, our study sought to compare two main forms of landscape matrix in western Rwanda: (1) tree plantations largely established through the recent FLR initiative, and (2) home gardens managed by local communities through long-standing traditional practices. We assessed how each type of landscape matrix influences landscape connectivity, and how these, in return, affect biodiversity. We chose woody plants and birds as biodiversity indicators, since they are among the most-studied and data-rich taxa (Stevenson and Fanshawe 2002; Fischer and Killmann 2008; Vande weghe and Vande weghe 2010; Fischer et al. 2024). Specifically, we (i) examined the spatial patterns of tree-cover connectivity versus home-garden connectivity, and (ii) we explored the relationships between connectivity and species richness (and diversity) of woody plants and birds.

## Methods

### Study area

Our study was carried out in the Western Province of Rwanda (hereafter referred to as western Rwanda), encompassing seven administrative districts, namely Karongi, Rutsiro, Rubavu, Nyabihu, Ngororero, Rusizi, Nyamasheke (Fig. 1). Western Rwanda covers an area of 5,877 km<sup>2</sup> on both sides of the Congo-Nile Crest (approx. 3,700 km<sup>2</sup>; Clay and Dejaeger 1987; Ndayambaje et al. 2014; Mukuralinda et al. 2016). The area is part of the Albertine Rift Ecoregion. Once characterized by a continuous forest belt, it has shrunk



**Fig. 1** Study area in the Western Province of Rwanda, covering seven administrative districts from north to south: Rubavu, Nyabihu, Rutsiro, Ngororero, Karongi, Nyamasheke

and Rusizi. The blue dotted line marks the Congo-Nile watershed divide (Congo-Nile Crest), while protected national park areas are highlighted in green color

to a few protected areas, namely Volcanoes National Park (NP), Nyungwe NP, and Gishwati-Mukura NP (UNESCO biosphere reserve since 2020; comprising Gishwati Forest and Mukura Forest; Republic of Rwanda 2020). Despite large parts of previously forested land being cleared, this montane forest ecoregion still provides vital habitat for a diverse biota with a high degree of endemism (Plumptre et al. 2007; Kindt 2014). Elevation in the study area ranges from 2,000 to 4,500 m a.s.l. (Mukashema et al. 2014), with a mean annual precipitation of about 2,000 mm

(Vande weghe and Vande weghe 2010), and an average annual mean temperature of 19 °C (World bank 2021).

The land-use history in western Rwanda was largely shaped by political events and socio-economic pressures (Akinyemi 2017), with an estimated 50 to 74% of natural Afromontane forest cleared for subsistence agriculture, livestock grazing, fuelwood, and construction materials (Kanyambwa 1998; Akinyemi 2017). Today, the landscape matrix in western Rwanda comprises different forms of land

use, namely tree plantations (e.g., *Eucalyptus*, *Grevillea*, *Pinus*, and *Alnus* species), subsistence farming (including small fields and home gardens), cash crop plantations (mainly tea), and pastures (Mukuralinda et al. 2016; Arakwiye et al. 2021). This complex mosaic reflects not only the historical land use patterns but also recent restoration interventions. For instance, in 2014, the government of Rwanda launched the Landscape Approach to Forest Restoration and Conservation (LAFREC) project, as part of the broader FLR initiative, and focused on restoring the Gishwati-Mukura landscape, a critical biodiversity hotspot within Rutsiro District (Republic of Rwanda 2025). To date, Rutsiro District comprises the largest area of tree plantations in Western Province, with FLR efforts adding about 818 hectares of new tree cover each year (Ministry of Environment 2019).

#### Assessing landscape connectivity across tree cover and home gardens

Landscape connectivity is defined as the degree to which a landscape facilitates or impedes movement among resource patches (Taylor et al. 1993), and is critical for supporting species movement, gene flow, and long-term biodiversity persistence at the landscape scale (Noss 1991; Rudnick et al. 2012). Connectivity can be conceptualized from multiple perspectives, including structural connectivity (the physical arrangement of vegetation/land-cover elements), functional or habitat connectivity (species-specific responses to landscape structure), and broader ecological connectivity (Lindenmayer and Fischer 2006). For this study, landscape connectivity was defined as the structural connectivity of vegetation/land cover, in our case, with a focus on trees and home gardens. We applied the circuit theory approach (McRae et al. 2016) to assess landscape connectivity across tree cover (here intentionally used as a highly simplified proxy that combines natural forests and tree plantations; Ministry of Environment 2019), and across home gardens. We used a set of openly available satellite-based datasets (see below) on tree cover and building structures. To implement the approach, we used the tool Omniscape which simulates electric current flow across the landscape (McRae et al. 2008, 2016; Dickson et al. 2019). The process starts by identifying the landscape type to be connected

and assigning resistance values to landscape features. A moving window procedure then connects all pixels within a given search radius, and the results from different window sizes are aggregated to generate a cumulative current map representing connectivity (McRae et al. 2016).

To map connectivity with a focus on tree cover, we used the ESA World Cover dataset 2021, which provides land-cover classes at a high pixel resolution of 10 m (Zanaga et al. 2022). In a first step, we masked out all classes other than trees. Second, we aggregated the remaining tree cover to a 100-m grid by counting the number of pixels containing trees in each grid cell. This yielded a fractional tree-cover map of trees at a 100-m spatial resolution. Third, we inverted this fractional cover dataset to assign lower resistance values (i.e., 0) to areas of relatively high tree cover, and high resistance values (i.e., 3; Phillips et al. 2021) to areas of low tree cover (e.g., Lake Kivu). We then set moving window radius of 1 km as the cutoff to assess tree-cover connectivity. This buffer was chosen because most birds have an activity radius exceeding 1 km, and they act as seed dispersers for indigenous trees—potentially disseminating seeds over even greater distances (Pejchar et al. 2008; Mueller et al. 2014).

To map connectivity across home gardens, we first identified home gardens in our study area based on Google Open Building polygon datasets collected in May 2023 (Sirko et al. 2021). This dataset contains different types of buildings, which are almost exclusively related to housing structures. Home gardens were operationally defined as the areas within a 15 m radius of each building. This buffer distance was chosen based on field knowledge of the study system, where most households practice subsistence agriculture and typically maintain small plots with woody vegetation—particularly fruit trees—immediately surrounding their homes. Similarly to the tree-cover mapping, we then calculated for each 100-m grid cell the fractional area covered with home gardens and inverted the resulting layer to generate a resistance layer, such that areas with many home gardens show lower resistance values, whereas areas with few home gardens show higher resistance values. Again, a radius of 1 km was assigned to assess the home-garden connectivity. To facilitate interpretation, both connectivity outputs (current flow) were transformed to a

scale from 1 to 100, where higher current flow values indicate stronger connectivity, and lower values represent weak or no connectivity. Prior to the modelling process, we applied a 30 km buffer around the study area—equivalent to 50% of the study area's width—to minimize edge effects and reduce bias (Koen et al. 2014; Pelletier et al. 2014). Omniscap modelling was conducted using the 'Omniscap.jl' software package within the 'Julia' programming environment version 0.6.2 (Landau et al. 2021).

### Relating biodiversity metrics to landscape connectivity

Once connectivity landscapes were established, we assessed how tree-cover and home-garden connectivity related to woody plant and bird richness and diversity, respectively. We established woody plant and bird richness and diversity at 91 sampling sites, including 47 tree plantations and 44 home gardens. Tree plantations were only sampled if the area exceeded one hectare and if they were not logged for more than one year. We focused on large plantation patches because they contribute more to landscape tree-cover connectivity and are expected to have stronger effects on biodiversity. Smaller patches vary widely in size and are more influenced by edge effects and disturbance, making consistent landscape-level comparisons difficult. We randomly chose home garden sites across the landscape to minimize spatial autocorrelation but accounted for road accessibility. Each site was sampled twice, once during the long dry season (20 June to 31 July 2024) and once during the short dry season (12 February to 5 March 2025). Woody plant richness was sampled by recording the presence and abundance of all woody plant species within a 20×20 m sampling plot at each site. Woody plants were defined as trees and shrubs growing taller than 1.3 m, identified using Fischer and Killmann (2008) and Fischer et al. (2024). For bird surveys, we applied the point count method (Ralph et al. 1995), i.e., standing for 15 min at the centre of each sampling plot to record and identify all birds seen or heard from any direction. Bird species were identified using Stevenson and Fanshawe (2002), Vande weghe and Vande weghe (2010) and Billerman et al. (2025). We conducted bird surveys only during days without rain or strong wind from 5:30 to 10:00 am. Based on habitat use and their functional role in the


ecosystem, birds were classified into four categories of forest dependence, namely, forest specialists, forest generalists, forest visitors, and non-forest species following Bennun et al. (1996; Supplementary Material Table 2). Based on the richness and abundance data of woody plants and birds, we calculated the respective Shannon diversity index using the 'vegan' package in R (Oksanen et al. 2001), before applying Mann–Whitney U tests to assess pairwise differences in observed species richness and diversity (for woody plants and birds separately) between tree plantations and home gardens. Finally, we assessed the relationship between both richness and diversity with our connectivity indices using Spearman's rank correlations.

## Results

### Tree-cover and home-garden connectivity

We obtained maps of tree-cover and home-garden connectivity at 100-m pixel resolution (Figs. 2, 3). Across the tree-cover connectivity map, the highest current flow values, indicating areas of strongest connectivity, were concentrated within the protected areas of Nyungwe NP, Gishwati-Mukura NP, and the Volcanoes NP (Fig. 2a, c–e). Outside these protected areas, strong tree-cover connectivity was observed in the northern part of Gishwati Forest (Fig. 2a, b), an area that received extensive FLR efforts in the past decade. Moderately high connectivity was observed in many parts of the landscape matrix adjacent to protected areas, including the western regions of Gishwati Forest (Fig. 2c), Mukura Forest (Fig. 2d) and Nyungwe NP (Fig. 2e), and some areas southeast of Volcanoes NP (Fig. 2a).

In contrast, home-garden connectivity map exhibited a distinct structural pattern. Strong connectivity was observed in the northern part of Gishwati Forest, along the main road connecting Rubavu in the West and Musanze City in the East (Fig. 3b). Another corridor is apparent along the Lake Kivu shoreline stretching from Rubavu southward to Cyangugu, i.e., an area located west of Gishwati-Mukura and Nyungwe NP (Fig. 3c, e). Moreover, south-eastern part of Volcanoes NP had a moderately high home-garden connectivity (Fig. 3d).



**Fig. 2** Tree-cover connectivity across western Rwanda. Panel A was modelled using a 100-m resolution resistance layer, with the moving window radius parameter set to 1 km in Omniscape. Panels B–E provide zoomed-in views of selected areas, with panels B–D also showing biodiversity survey sites

located within tree plantations. The color gradient shows tree-cover connectivity percentiles for the landscape: red areas indicate high tree-cover connectivity, while blue areas highlight barriers with low or zero connectivity


A moderately high home-garden connectivity was observed across most parts of the landscape matrix.

#### Correlation between biodiversity metrics and landscape connectivity

Across the 47 tree plantation sites, we recorded eight woody plant species (7 exotic and 1 native species), with each site containing one to three species (Supplementary Table 1). In contrast, the 44 home garden sites supported a higher diversity of 24 woody plant species (18 exotic and 6 native species), with one to ten species per site (Supplementary Table 1). Furthermore, we recorded 32 bird species at tree plantations and 64 species in the home gardens (Supplementary Table 2). Most bird species observed were small-bodied forest visitors or non-forest species. All species

had a conservation status of ‘Least Concern’ (Billerman et al. 2025).

Mann-Whitney U test showed that both species richness and diversity of woody plants and birds differed highly significant between tree plantations and home gardens (woody plant richness:  $W = 1997$ ,  $p < 0.001$ ; woody plant diversity:  $W = 2068$ ,  $p < 0.001$ ; bird richness:  $W = 1606$ ,  $p < 0.001$ ; bird diversity:  $W = 1576$ ,  $p < 0.001$ ). Species richness (mean  $\pm$  SD) was significantly higher in home gardens (woody plants:  $4.16 \pm 1.95$ ; birds:  $4.84 \pm 4.00$ ) than at tree plantations (woody plants:  $1.34 \pm 0.52$ ; birds:  $1.81 \pm 1.75$ ; Fig. 4a, c). The Shannon diversity index (mean  $\pm$  SD) was also significantly higher in home gardens (woody plants:  $1.27 \pm 0.41$ ; birds:  $1.20 \pm 0.75$ ) than in tree plantations (woody plants:  $0.15 \pm 0.25$ ; birds:  $0.47 \pm 0.62$ ; Fig. 4b, d).




**Fig. 3** Panel A. Home-garden connectivity was modelled using a 100-m resolution resistance layer, with the moving window radius parameter set to 1 km in Omniscape. Panels B–E provide zoomed-in views of areas with relatively high home-garden connectivity. The color gradient represents

home-garden connectivity percentiles across the landscape: red areas indicate high home-garden connectivity, while blue areas highlight barriers with low or zero connectivity. White circles show biodiversity survey sites in home gardens

Spearman's rank correlations revealed a significant negative correlation between tree-cover connectivity and home-garden connectivity ( $r = -0.71$ ,  $p < 0.001$ , Fig. 5). In general, we observed a higher richness and diversity of woody plants (Fig. 5a, b) and birds (Fig. 5c, d) in home gardens than in tree plantations. Highest bird richness and diversity (Fig. 5c, d) clustered in areas of medium to high home-garden connectivity, associated with low to medium tree-cover connectivity. Conversely, areas with the highest tree-cover connectivity exhibited very low woody plant and bird richness and diversity.

Moreover, Spearman's rank correlations revealed a significant negative relationship between tree-cover connectivity and woody plant richness ( $r = -0.63$ ,  $p < 0.001$ ), as well as between tree-cover connectivity and bird richness ( $r = -0.46$ ,  $p < 0.001$ ). We also found

a significant negative correlation between tree-cover connectivity and woody plant diversity ( $r = -0.67$ ,  $p < 0.001$ ) and between tree-cover connectivity and bird diversity ( $r = -0.45$ ,  $p < 0.001$ ). In contrast, we found a significant positive relationship between home-garden connectivity and woody plant richness ( $r = 0.64$ ,  $p < 0.001$ ), as well as between home-garden connectivity and bird richness ( $r = 0.44$ ,  $p < 0.001$ ). Our analysis further revealed a significant positive correlation between home-garden connectivity and woody plant diversity ( $r = 0.68$ ,  $p < 0.001$ ), and between home-garden connectivity and bird diversity ( $r = 0.41$ ,  $p < 0.001$ ).




**Fig. 4** Boxplots showing the species richness and diversity of woody plants and birds recorded in tree plantations (black color) and home gardens (grey color): **A** woody plant richness; **B** Shannon index of woody plant diversity; **C** bird

species richness; **D** Shannon index of bird diversity. Results of pairwise comparisons using Mann–Whitney U tests are indicated within each plot

## Discussion

Many regions in the tropics have suffered from considerable forest loss, fragmentation, and degradation in the past, and tree planting efforts seek to reverse this. How such tree plantations contribute to this goal, and what their value is relative to other land uses that contain tree cover, such as home gardens, remains poorly understood. Here, in a global

hotspot of tropical forest restoration, we assessed connectivity of two different forms of land use (i.e., recently established tree plantations and traditional home gardens) in relation to two biodiversity metrics (i.e., woody plant and bird richness and diversity). By comparing connectivity maps for both forms of land use, and by relating connectivity to diversity metrics, we found tree plantations to function like a ‘hard matrix’, offering only limited habitat value and



**Fig. 5** Scatterplots showing the relationships between tree-cover connectivity and home-garden connectivity, with **A** woody plant richness; **B** Shannon diversity index of woody plants; **C** bird richness; **D** Shannon diversity index of birds

represented by a color gradient ranging from blue (low richness) to red (high richness). Points represent different land matrix types: circles for home gardens and triangles for tree plantations

restricted permeability, while home gardens serve as a ‘soft matrix’ allowing organisms to disperse (sensu Franklin 1993; Lindenmayer and Fischer 2006). Our findings challenge conventional tree plantations in FLR practice and underscore the need

to actively incorporate home gardens in restoration strategies. We discuss our findings in the context of two implications for FLR policy and practice: (i) tree plantations increase tree-cover connectivity but not biodiversity, and (ii) home gardens may serve as

stepping stones for species dispersal and thus benefit biodiversity conservation.

Tree plantations increase tree-cover connectivity but not biodiversity

FLR practice has contributed to achieving Rwanda's national target of 30% forest cover, as outlined in the Vision 2020 (Republic of Rwanda 2012). Our analyses indicate that FLR-led restoration efforts have successfully increased tree cover in the landscape, leading to high tree-cover connectivity around protected areas (Fig. 2a-e). However, well-connected areas of trees supported significantly lower richness and diversity of woody plants and birds than less well-connected areas (Fig. 5a-d). As such, it appears that tree planting efforts to date have increased connectivity but failed to enhance habitat quality and thus biodiversity. Compared to previous studies (e.g., Crouzeilles et al. 2019) that identified tree-cover and tree-cover connectivity as key indicators for the recovery of tropical biodiversity (and as indicators of restoration success), our findings offer a contrasting view.

Sustainably managed tree plantations that do not replace natural or semi-natural ecosystems could contribute to the conservation of native biodiversity, generally hosting fewer species than natural forests (Hua et al. 2022; Wang et al. 2022), but more than other human-modified land use types (Brockerhoff et al. 2008). For example, a study in Uganda's Kibale NP, showed that planting native tree species can accelerate biodiversity recovery within a relatively short period (Omeja et al. 2011). In a densely populated landscape like western Rwanda, tree plantations can play an indirect, socially important buffering role by supplying wood and reducing pressure on natural forests. Restoring landscapes with indigenous species instead of exotic plantations could help mitigate anthropogenic pressures on natural forests through community-based forest management, sustainable use zones that allow regulated harvesting, and alternative energy and livelihood options to reduce fuelwood dependence.

FLR-led restoration efforts have primarily been achieved by the widespread establishment of exotic tree plantations, now comprising more than half of the country's forested area (Ministry of Environment 2019). Our findings suggest that, at least for woody

plants and birds, these plantations—especially when disconnected from natural forest patches—function as a 'hard matrix' with limited habitat value and low attractiveness to biodiversity. From the landscape perspective, if this type of restoration practice continues, the landscape matrix may facilitate species invasiveness, eventually leading to biotic homogenization, where diverse native species are replaced by a few widespread, fast-growing exotics, consequently increasing the extinction risk for native species (Crosby 2004; Holl et al. 2022). Such homogenization is already evident in our data, as bird communities in home gardens and forest plantations showed substantial overlap. We thus recommend FLR practice in western Rwanda and other regions of the Albertine Rift to shift from overly simple, short-term, target-based indicators such as tree cover towards strategies that reflect long-term restoration success and ecological functionality, such as the promotion of native vegetation (Raj et al. 2023) and the creation of a mosaic of resources and habitat features that facilitate animal dispersal, foraging, and reproduction (Holl et al. 2022).

Home gardens could serve as stepping stones for biodiversity

Recent studies showed home gardens in Rwanda are linked to improved conservation of wild plants (Nsengimana et al. 2020), to increased medicinal plant diversity (Rizinjirabake et al. 2024), and to improvements in dietary diversity and thus higher nutritional value of local communities (Issahaku et al. 2023; Dusingizimana et al. 2025). Until to date, little is known about the role of home gardens in supporting connectivity and biodiversity within the landscape matrix. Our study addressed this knowledge gap and showed that home gardens maintain intermediary habitats, following natural forests, in supporting relatively high richness and diversity of woody plants and birds. This is particularly valuable for connectivity since home gardens follow linear features in the landscape, such as along tracks and roads. Thus, our findings suggest that spatially interconnected home gardens may function as a 'soft matrix', positively contributing to biodiversity, even at locations where tree-cover connectivity is low. Unlike tree plantations, home gardens are more heterogeneous and offer ecologically valuable habitats, including indigenous

trees, shrubs and herbaceous plants arranged in a multistorey system (Hylander and Nemomissa 2009; Galluzzi et al. 2010), thereby attracting a variety of bird species (Engelen et al. 2017; Vargas-Cárdenas et al. 2024). While home gardens offer important benefits, they cannot substitute for natural forests. Our findings indicate that bird communities in home gardens are largely composed of forest visitors and non-forest species predominantly classified as ‘Least Concern’, suggesting that home gardens cannot replace natural forests in providing essential habitat for forest-dependent specialists. Similar patterns have been reported in the fragmented landscapes of Bangladesh (Bardhan et al. 2012), Guatemala (Haggar et al. 2019), and Ethiopia (Tesfay et al. 2025), where home-garden agroforestry systems function as stepping stones that enhance biodiversity conservation, acting as vital intermediary habitats that bridge and buffer natural forests within the broader landscape.

Agroforestry, including multi-layered home gardens, has been widely recognized as a key restoration element in Rwanda (Ministry of Natural Resources 2014). However, past restoration practices have primarily been based on the ‘tree-based approach’, which integrates trees with crops to improve soil fertility and water quality (Ministry of Natural Resources 2014; Iiyama et al. 2018). This practice has led to the widespread use of exotic *Grevillea robusta*, which provides little value for native biodiversity (Ruticugambi et al. 2024). Traditional home gardens have received relatively little attention in conservation and restoration efforts to date. This is due to an ever-increasing mechanisation in agriculture, the use of pesticides, and an increasing number of exotic plants. Such modernization processes make traditional home gardens face great challenges, eventually resulting in a sharp decline in species richness of indigenous plants, insects and birds (Seburanga 2013). Strengthening the conservation of traditional home gardens is therefore critical to sustaining their biodiversity value and ensuring they remain vital components of future restoration landscapes.

## Conclusion

Forest loss, fragmentation, and degradation are major drivers of biodiversity loss in tropical landscapes. While conserving natural forests

is paramount, scaling up forest restoration to improve degraded forests and connectivity among remaining patches through raising matrix quality is also vital for biodiversity conservation. Based on our findings, we recommend that restoration efforts prioritize the protection of remaining natural forests, which are irreplaceable for sustaining native biodiversity. Complementary strategies should promote the regeneration of secondary forests dominated by native tree taxa and the expansion of biodiverse agroforestry systems. Traditional home gardens, integrating native plant species and traditional crops, offer a promising social–ecological restoration approach. In western Rwanda, the comparison of two tree-rich land-use systems showed that exotic tree plantations, despite enhancing tree-cover connectivity, provide limited biodiversity benefits, whereas traditional home gardens function as valuable ecological stepping stones. In conclusion, to maximize forest landscape restoration effectiveness in biodiversity-rich regions, we recommend promoting native tree plantations alongside biodiversity-friendly, tree-rich systems such as home gardens.

**Acknowledgements** We gratefully acknowledge funding by the Deutsche Forschungsgemeinschaft (DFG) — project number 496337053. We also extend our thanks to the National Council for Science and Technology of Rwanda for authorizing the research permit in Rwanda (permit reference number: NCST/482/0148/2024). PS would like to thank Shimin Pan, Dominic Munyensanga, Flugence Iyamuremye, and Jean Damascene Bariyanga for their invaluable assistance with fieldwork. We are grateful to three anonymous reviewers, whose comments and constructive feedback has helped us to improve this manuscript.

**Author contributions** PS, MB, JF conceptualisation; JF and TK funding acquisition; PS and MB spatial data collection; PS field data collection; PS data curation, data analysis; PS wrote original draft; MB, JF, BK, TK supervision. All authors reviewed and edited the manuscript.

**Funding** Open Access funding enabled and organized by Projekt DEAL. This work was supported by the Deutsche Forschungsgemeinschaft (DFG) – project number 496337053, as part of the research unit “A social-ecological systems approach to inform ecosystem restoration in rural Africa”. This publication was funded by the Open Access Publication Fund of Leuphana University Lüneburg.

**Data availability** All data supporting the findings of this study are provided within the paper and its Supplementary Material and Supplementary Data.

## Declarations

**Competing interest** The authors have no relevant financial or non-financial interests to disclose. The authors declare no competing interests.

**Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

## References

- Abebe T, Sterck FJ, Wiersum KF, Bongers F (2013) Diversity, composition and density of trees and shrubs in agroforestry homegardens in Southern Ethiopia. *Agrofor Syst* 87:1283–1293. <https://doi.org/10.1007/s10457-013-9637-6>
- Akinyemi FO (2017) Land change in the central Albertine Rift: insights from analysis and mapping of land use-land cover change in north-western Rwanda. *Appl Geogr* 87:127–138. <https://doi.org/10.1016/j.apgeog.2017.07.016>
- Anderson J, Rowcliffe JM, Cowlshaw G (2007) Does the matrix matter? A forest primate in a complex agricultural landscape. *Biol Conserv* 135:212–222. <https://doi.org/10.1016/j.biocon.2006.10.022>
- Arakwiye B, Rogan J, Eastman JR (2021) Thirty years of forest-cover change in Western Rwanda during periods of wars and environmental policy shifts. *Reg Environ Change* 21:27. <https://doi.org/10.1007/s10113-020-01744-0>
- Bardhan S, Jose S, Biswas S et al (2012) Homegarden agroforestry systems: an intermediary for biodiversity conservation in Bangladesh. *Agrofor Syst* 85:29–34. <https://doi.org/10.1007/s10457-012-9515-7>
- Bennun L, Dranzoa C, Pomeroy D (1996) The forest birds of Kenya and Uganda. *J East Afr Nat Hist* 85:23–48. [https://doi.org/10.2982/0012-8317\(1996\)85\[23:TFBOKA\]2.0.CO;2](https://doi.org/10.2982/0012-8317(1996)85[23:TFBOKA]2.0.CO;2)
- Bhagwat SA, Willis KJ, Birks HJB et al (2008) Agroforestry: a refuge for tropical biodiversity? *Trends Ecol Evol* 23(5):261–267. <https://doi.org/10.1016/j.tree.2008.01.005>
- Billerman SM, Keeney BK, Kirwan GM et al (2025) Birds of the World. <https://birdsoftheworld.org/bow/home> Accessed 12 June 2025
- Boesing AL, Nichols E, Metzger JP (2018) Biodiversity extinction thresholds are modulated by matrix type. *Ecography* 41:1520–1533. <https://doi.org/10.1111/ecog.03365>
- Brançalion PHS, Chazdon RL (2017) Beyond hectares: four principles to guide reforestation in the context of tropical forest and landscape restoration. *Restor Ecol* 25:491–496. <https://doi.org/10.1111/rec.12519>
- Brinck K, Fischer R, Groeneveld J, Lehmann S, Dantas De Paula M, Pütz S, Sexton JO, Song D, Huth A (2017) High resolution analysis of tropical forest fragmentation and its impact on the global carbon cycle. *Nat Commun* 8:14855. <https://doi.org/10.1038/ncomms14855>
- Brockerhoff EG, Jactel H, Parrotta JA et al (2008) Plantation forests and biodiversity: oxymoron or opportunity? *Biodivers Conserv* 17:925–995. <https://doi.org/10.1007/s10531-008-9380-x>
- Bullock JM, Aronson J, Newton AC, Pywell RF, Rey-Benayas JM (2011) Restoration of ecosystem services and biodiversity: conflicts and opportunities. *Trends Ecol Evol* 26:541–549. <https://doi.org/10.1016/j.tree.2011.06.011>
- Chazdon RL, Guariguata MR (2018) Decision support tools for forest landscape restoration: current status and future outlook. *Cent Int For Res (CIFOR)*. <https://doi.org/10.17528/cifor/006792>
- Chazdon RL, Harvey CA, Komar O, Griffith DM, Ferguson BG, Martínez-Ramos M, Morales H, Nigh R, Soto-Pinto L, Van Breugel M, Philpott SM (2009) Beyond reserves: a research agenda for conserving biodiversity in human-modified tropical landscapes. *Biotropica* 41:142–153. <https://doi.org/10.1111/j.1744-7429.2008.00471.x>
- Chazdon RL, Brancalion PHS, Lamb D, Laestadius L, Calmon M, Kumar C (2017) A policy-driven knowledge agenda for global forest and landscape restoration. *Conserv Lett* 10:125–132. <https://doi.org/10.1111/conl.12220>
- Chazdon RL, Gutierrez V, Brancalion PHS et al (2020) Co-creating conceptual and working frameworks for implementing forest and landscape restoration based on core principles. *Forests* 11(6):706. <https://doi.org/10.3390/f11060706>
- Clay DC, Dejaegher YMJ (1987) Agro-ecological zones: the development of a regional classification scheme for Rwanda. *Tropicultura* 5:153–159
- Clewell A, Aronson J, Winterhalder K (2004) The SER international primer on ecological restoration. Society for Ecological Restoration International Science & Policy Working Group. <https://www.ser.org/page/SERDocuments> Accessed 10 June 2025
- Crosby A (2004) Ecological imperialism: The biological expansion of Europe, 900–1900, 2nd edn. Cambridge University Press
- Crouzeilles R, Barros FSM, Molin PG, Ferreira MS, Junqueira AB, Chazdon RL, Lindenmayer DB, Tymus JRC, Strassburg BBN, Brancalion PHS (2019) A new approach to map landscape variation in forest restoration success in tropical and temperate forest biomes. *J Appl Ecol* 56:2675–2686. <https://doi.org/10.1111/1365-2664.13501>
- de Souza Leite M, Boesing AL, Metzger JP, Prado PI (2022) Matrix quality determines the strength of habitat loss

- filtering on bird communities at the landscape scale. *J Appl Ecol* 59:2790–2802. <https://doi.org/10.1111/1365-2664.14275>
- Dickson BG, Albano CM, Anantharaman R, Beier P, Fargione J, Graves TA, Gray ME, Hall KR, Lawler JJ, Leonard PB, Littlefield CE, McClure ML, Novembre J, Schloss CA, Schumaker NH, Shah VB, Theobald DM (2019) Circuit-theory applications to connectivity science and conservation. *Conserv Biol* 33:239–249. <https://doi.org/10.1111/cobi.13230>
- Dobrovolski R, Diniz-Filho JAF, Loyola RD, De Marco P (2011) Agricultural expansion and the fate of global conservation priorities. *Biodivers Conserv* 20:2445–2459. <https://doi.org/10.1007/s10531-011-9997-z>
- Dusingizimana T, Nduwayezu G, Kjelqvist T (2025) Women's dietary diversity is associated with homestead production and market access: a cross-sectional study in rural Rwanda. *Matern Child Nutr* 21:e13755. <https://doi.org/10.1111/mcn.13755>
- Ellis EC, Gauthier N, Klein Goldewijk K et al (2021) People have shaped most of terrestrial nature for at least 12,000 years. *PNAS* 118:e2023483118. <https://doi.org/10.1073/pnas.2023483118>
- Engelen D, Lemessa D, Şekerciöğlü ÇH, Hylander K (2017) Similar bird communities in homegardens at different distances from Afromontane forests. *Bird Conserv Int* 27:83–95. <https://doi.org/10.1017/S0959270916000162>
- Erdmann TK (2005) Agroforestry as a tool for restoring forest landscapes. In: Mansourian S, Vallauri D, Dudley N (eds) *Forest restoration in landscapes*. Springer, New York, pp 274–284
- Fahrig L (2003) Effects of habitat fragmentation on biodiversity. *Annu Rev Ecol Evol Syst* 34:487–515. <https://doi.org/10.1146/annurev.ecolsys.34.011802.132419>
- FAO (2017) *Agroforestry for landscape restoration: exploring the potential of agroforestry to enhance the sustainability and resilience of degraded landscapes*. Rome. <https://doi.org/10.4060/i7374e>
- Fischer E, Killmann D (2008) *Illustrated field guide to the plants of Nyungwe National Park Rwanda*. University of Koblenz-Landau, Koblenz
- Fischer J, Lindenmayer DB (2007) Landscape modification and habitat fragmentation: a synthesis. *Glob Ecol Biogeogr* 16:265–280. <https://doi.org/10.1111/j.1466-8238.2007.00287.x>
- Fischer E, Nsanzurwimo A, Dumbo B, Richter R, Weghe JPV (2024) An updated checklist of vascular plants (lycophytes, ferns, gymnosperms, angiosperms) from Nyungwe National Park (incl. Cyamudongo Forest), Rwanda. *Phytotaxa* 673:1–113. <https://doi.org/10.11646/phytotaxa.673.1.1>
- Franklin JF (1993) Preserving biodiversity: species, ecosystems, or landscapes? *Ecol Appl* 3:202–205. <https://doi.org/10.2307/1941820>
- Galhena DH, Freed R, Maredia KM (2013) Home gardens: a promising approach to enhance household food security and wellbeing. *Agric Food Secur* 2:8. <https://doi.org/10.1186/2048-7010-2-8>
- Galluzzi G, Eyzaguirre P, Negri V (2010) Home gardens: neglected hotspots of agro-biodiversity and cultural diversity. *Biodivers Conserv* 19:3635–3654. <https://doi.org/10.1007/s10531-010-9919-5>
- Gascon C, Lovejoy TE, Bierregaard RO Jr et al (1999) Matrix habitat and species richness in tropical forest remnants. *Biol Conserv* 91(2–3):223–229. [https://doi.org/10.1016/s0006-3207\(99\)00080-4](https://doi.org/10.1016/s0006-3207(99)00080-4)
- Gbedomon RC, Salako VK, Fandohan AB, Idohou AFR, Glèlè Kakai R, Assogbadjo AE (2017) Functional diversity of home gardens and their agrobiodiversity conservation benefits in Benin, West Africa. *J Ethnobiol Ethnomed* 13:66. <https://doi.org/10.1186/s13002-017-0192-5>
- Gibson L, Lee T, Koh L et al (2021) Primary forests are irreplaceable for sustaining tropical biodiversity. *Nature* 478:378–381. <https://doi.org/10.1038/nature10425>
- Gifawesen ST, Tola FK, Duguma MS (2020) Review on role of home garden agroforestry practices to improve livelihood of small scale farmers and climate change adaptation and mitigation. *J Plant Sci* 8(5):134–145. <https://doi.org/10.11648/j.jps.20200805.15>
- Gonzalez A, Mouquet N, Loreau M (2009) Biodiversity as spatial insurance: The effects of habitat fragmentation and dispersal on ecosystem functioning. In: Naem S, Bunker DE, Hector A, Loreau M, Perrings C (eds) *Biodiversity, ecosystem functioning and ecosystem services*. Oxford University Press, Oxford, pp 134–146
- Haddad NM, Brudvig LA, Clobert J, Davies KF et al (2015) Habitat fragmentation and its lasting impact on Earth's ecosystems. *Sci Adv* 1:e1500052. <https://doi.org/10.1126/sciadv.1500052>
- Hagen M, Kissling WD, Rasmussen C et al (2012) Biodiversity, species interactions and ecological networks in a fragmented world. *Adv Ecol Res* 46:89–210. <https://doi.org/10.1016/B978-0-12-396992-7.00002-2>
- Haggard J, Pons D, Saenz L, Vides M (2019) Contribution of agroforestry systems to sustaining biodiversity in fragmented forest landscapes. *Agric Ecosyst Environ* 283:106567. <https://doi.org/10.1016/j.agee.2019.06.006>
- Hansen L, Sorgho R, Mank I, Nayna Schwerdtle P, Agure E, Bärnighausen T, Danquah I (2022) Home gardening in sub-Saharan Africa: a scoping review on practices and nutrition outcomes in rural Burkina Faso and Kenya. *Food Energy Secur* 11:e388. <https://doi.org/10.1002/fes3.388>
- Hatfield JH, Harrison MLK, Banks-Leite C (2018) Functional diversity metrics: how they are affected by landscape change and how they represent ecosystem functioning in the tropics. *Curr Landsc Ecol Rep* 3:35–42. <https://doi.org/10.1007/s40823-018-0032-x>
- Holl KD, Luong JC, Brancalion PHS (2022) Overcoming biotic homogenization in ecological restoration. *Trends Ecol Evol* 37:777–788. <https://doi.org/10.1016/j.tree.2022.05.002>
- Hua F, Bruijnzeel LA, Meli P et al (2022) The biodiversity and ecosystem service contributions and trade-offs of forest restoration approaches. *Science* 376:839–844. <https://doi.org/10.1126/science.abl4649>
- Hylander K, Nemomissa S (2009) Complementary roles of home gardens and exotic tree plantations as alternative habitats for plants of the Ethiopian montane rainforest.

- Conserv Biol 23:400–409. <https://doi.org/10.1111/j.1523-1739.2008.01097.x>
- Iiyama M, Mukuralinda A, Ndayambaje JD, Musana B, Ndoli A, Mowo JG, Garrity D, Ling S, Ruganzu V (2018) Tree-based ecosystem approaches (TBEAs) as multi-functional land management strategies—evidence from Rwanda. *Sustainability* 10:1360. <https://doi.org/10.3390/su10051360>
- Issahaku G, Kornher L, Saiful Islam AHM, Abdul-Rahaman A (2023) Heterogeneous impacts of home-gardening on household food and nutrition security in Rwanda. *Food Secur* 15:731–750. <https://doi.org/10.1007/s12571-023-01344-w>
- Jenkins CN, Pimm SL, Joppa LN (2013) Global patterns of terrestrial vertebrate diversity and conservation. *Proc Natl Acad Sci U S A* 110:e2602–e2610. <https://doi.org/10.1073/pnas.130225111>
- Jules ES, Shahani P (2003) A broader ecological context to habitat fragmentation: why matrix habitat is more important than we thought. *J Veg Sci* 14(3):459–464. <https://doi.org/10.1111/j.1654-1103.2003.tb02172.x>
- Kanyamibwa S (1998) Impact of war on conservation: Rwandan environment and wildlife in agony. *Biodivers Conserv* 7:1399–1406. <https://doi.org/10.1023/A:1008880113990>
- Kindt R (2014) Potential natural vegetation of Eastern Africa (Ethiopia, Kenya, Malawi, Rwanda, Tanzania, Uganda and Zambia). University of Copenhagen, Department of Geoscience and Natural Resource Management
- Koen EL, Bowman J, Sadowski C, Walpole AA (2014) Landscape connectivity for wildlife: development and validation of multispecies linkage maps. *Methods Ecol Evol* 5(7):626–633. <https://doi.org/10.1111/2041-210X.12197>
- Kremen C, Merenlender AM (2018) Landscapes that work for biodiversity and people. *Science* 362:eaa6020. <https://doi.org/10.1126/science.aau6020>
- Kumar BM, Nair PKR (eds) (2006) *Tropical homegardens: a time-tested example of sustainable agroforestry*. Springer, Netherlands
- Kupfer JA, Malanson GP, Franklin SB (2006) Not seeing the ocean for the islands: the mediating influence of matrix-based processes on forest fragmentation effects. *Glob Ecol Biogeogr* 15:8–20. <https://doi.org/10.1111/j.1466-822X.2006.00204.x>
- Lamb D (2014) *Large-scale forest restoration*. Routledge. <https://doi.org/10.4324/9780203071649>
- Landau V, Shah V, Anantharaman R, Hall K (2021) Omniscap.jl: software to compute omnidirectional landscape connectivity. *J Open Source Softw* 6:2829. <https://doi.org/10.21105/joss.02829>
- Lewis SL, Edwards DP, Galbraith D (2015) Increasing human dominance of tropical forests. *Science* 349:827–832. <https://doi.org/10.1126/science.aaa9932>
- Lindenmayer DB, Fischer J (2006) *Landscape change and habitat fragmentation*. Island Press
- López-Bedoya PA, Magura T, Edwards FA et al (2021) What level of native beetle diversity can be supported by forestry plantations? A global synthesis. *Insect Conserv Divers* 14:736–747. <https://doi.org/10.1111/icad.12518>
- Maginnis S, Rietbergen-McCracken J, Sarre A (eds) (2012) *The forest landscape restoration handbook*. Routledge
- Mansourian S, Berrahmouni N (2021) *Review of forest and landscape restoration in Africa 2021*. FAO and AUDA-NEPAD. <https://doi.org/10.4060/cb6111en>
- Mansourian S, Stanturf JA, Derkyi MAA, Engel VL (2017) Forest landscape restoration: increasing the positive impacts of forest restoration or simply the area under tree cover? *Restor Ecol* 25:178–183. <https://doi.org/10.1111/rec.12489>
- Maroyi A (2009) Traditional homegardens and rural livelihoods in Nhema, Zimbabwe: a sustainable agroforestry system. *Int J Sustain Dev World Ecol* 16:1–8. <https://doi.org/10.1080/13504500902745895>
- McRae BH, Dickson BG, Keitt TH, Shah VB (2008) Using circuit theory to model connectivity in ecology, evolution, and conservation. *Ecology* 89:2712–2724. <https://doi.org/10.1890/07-1861.1>
- McRae B, Popper K, Jones A, Schindel M, Buttrick S, Hall K, Unnasch RS, Platt J (2016) Conserving nature’s stage: mapping omnidirectional connectivity for resilient terrestrial landscapes in the Pacific Northwest. The Nature Conservancy, Portland, Oregon
- Ministry of Environment (2019) *Rwanda Forest Cover Mapping*. Republic of Rwanda. [https://www.environment.gov.rw/fileadmin/user\\_upload/Moe/Publications/Reports/Forest\\_cover\\_report\\_2019.pdf](https://www.environment.gov.rw/fileadmin/user_upload/Moe/Publications/Reports/Forest_cover_report_2019.pdf) Accessed 12 June 2025
- Ministry of Natural Resources (2014) *Forest landscape restoration opportunity assessment for Rwanda*. MINIRENA (Rwanda), IUCN, WRI
- Mitchell MGE, Suarez-Castro AF, Martinez-Harms M, Maron M, McAlpine C, Gaston KJ, Johansen K, Rhodes JR (2015) Reframing landscape fragmentation’s effects on ecosystem services. *Trends Ecol Evol* 30:190–198. <https://doi.org/10.1016/j.tree.2015.01.011>
- Mittermeier RA, Turner WR, Larsen FW et al (2011) Global biodiversity conservation: the critical role of hotspots. In: Zachos F, Habel J (eds) *Biodiversity hotspots*. Springer, Berlin, pp 3–22. [https://doi.org/10.1007/978-3-642-20992-5\\_1](https://doi.org/10.1007/978-3-642-20992-5_1)
- Montagnini F (2020) The contribution of agroforestry to restoration and conservation: biodiversity islands in degraded landscapes. In: Dagar JC, Gupta SR, Teketay D (eds) *Agroforestry for degraded landscapes*. Springer Singapore
- Mueller T, Lenz J, Caprano T, Fiedler W, Böhning-Gaese K (2014) Large frugivorous birds facilitate functional connectivity of fragmented landscapes. *J Appl Ecol* 51:684–692. <https://doi.org/10.1111/1365-2664.12247>
- Mugabowindekwe M, Brandt M, Chave J, Reiner F, Skole DL, Kariryaa A, Igel C, Hiernaux P, Ciaia P, Mertz O, Tong X, Li S, Rwanyiziri G, Dushimiyimana T, Ndoli A, Uwizeyimana V, Lillesø J-PB, Gieseke F, Tucker CJ, Fensholt R (2023) Nation-wide mapping of tree-level aboveground carbon stocks in Rwanda. *Nat Clim Chang* 13:91–97. <https://doi.org/10.1038/s41558-022-01544-w>
- Mukashema A, Veldkamp A, Vrieling A (2014) Automated high resolution mapping of coffee in Rwanda using an expert Bayesian network. *Int J Appl Earth Obs Geoinf* 33:331–340. <https://doi.org/10.1016/j.jag.2014.05.005>

- Mukuralinda A, Ndayambaje JD, Iiyama M et al (2016) Taking to scale tree-based systems in Rwanda to enhance food security, restore degraded land, improve resilience to climate change and sequester carbon. PROFOR. [https://www.profor.info/sites/default/files/TakingToScaleRwanda\\_low.pdf](https://www.profor.info/sites/default/files/TakingToScaleRwanda_low.pdf) Accessed 12 June 2025
- Myers N (1988) Threatened biotas: “hot spots” in tropical forests. *Environist* 8:187–208. <https://doi.org/10.1007/BF02240252>
- Nair PR, Kumar BM, Nair VD (2021) An introduction to agroforestry: Four decades of scientific developments, Springer.
- National Institute of Statistics of Rwanda (2023) Rwanda Statistics Yearbook 2023. <https://www.statistics.gov.rw/>
- Ndayambaje JD, Mugiraneza T, Mohren GMJ (2014) Woody biomass on farms and in the landscapes of Rwanda. *Agrofor Syst* 88:101–124. <https://doi.org/10.1007/s10457-013-9659-0>
- Nduwamungu J (2011) Forest plantations and woodlots in Rwanda. *Afr for Forum* 1:2–61
- Noss RF (1991) Landscape connectivity: Different functions at different scales. In: *Landscape linkage and biodiversity*, University of Florida
- Nsengimana T, Nsengimana V, Nsanganwimana F (2020) Local knowledge and use of wild edible plants in eastern part of Nyungwe National Park in Rwanda: prospects for forest biodiversity conservation. *J Res For Wildl Environ* 12:854
- Oksanen J, Simpson GL, Blanchet FG, Kindt R et al (2001) vegan: Community Ecology Package (p 2.6–8) . <https://doi.org/10.32614/CRAN.package.vegan>
- Omeja PA, Chapman CA, Obua J, Lwanga JS, Jacob AL, Wanyama F, Mugenyi R (2011) Intensive tree planting facilitates tropical forest biodiversity and biomass accumulation in Kibale National Park, Uganda. *For Ecol Manag* 261:703–709. <https://doi.org/10.1016/j.foreco.2010.11.029>
- Panyadee P, Balslev H, Wangpakapattanawong P, Inta A (2018) Karen homegardens: characteristics, functions, and species diversity. *Econ Bot* 72:1–19. <https://doi.org/10.1007/s12231-018-9404-8>
- Peh KSH, Lin YangChen LY, Luke SH, Foster WA, Turner EC (2014) Forest fragmentation and ecosystem function. In: Kettle CJ, Koh LP (eds) *Global forest fragmentation*, 1st edn. CABI
- Pejchar L, Pringle RM, Ranganathan J et al (2008) Birds as agents of seed dispersal in a human-dominated landscape in southern Costa Rica. *Biol Conserv* 141:536–544. <https://doi.org/10.1016/j.biocon.2007.11.008>
- Pelletier D, Clark M, Anderson MG et al (2014) Applying circuit theory for corridor expansion and management at regional scales: tiling, pinch points, and omnidirectional connectivity. *PLoS ONE* 9(1):e84135. <https://doi.org/10.1371/journal.pone.0084135>
- Perino A, Pereira HM, Navarro LM, Fernández N, Bullock JM, Ceaușu S, Cortés-Avизanda A, Van Klink R, Kuemmerle T, Lomba A, Pe'er G, Plieninger T, Rey Benayas JM, Sandom CJ, Svenning JC, Wheeler HC (2019) Rewilding complex ecosystems. *Science* 364:eav5570. <https://doi.org/10.1126/science.aav5570>
- Phillips P, Clark MM, Baral S et al (2021) Comparison of methods for estimating omnidirectional landscape connectivity. *Landsc Ecol* 36:1647–2166. <https://doi.org/10.1007/s10980-021-01254-2>
- Plumptre AJ, Davenport TRB, Behangana M, Kityo R, Eilu G, Ssegawa P, Ewango C, Meirte D, Kahindo C, Herremans M, Peterhans JK, Pilgrim JD, Wilson M, Languy M, Moyer D (2007) The biodiversity of the Albertine Rift. *Biol Conserv* 134:178–194. <https://doi.org/10.1016/j.biocon.2006.08.021>
- Raj A, Jhariya MK, Banerjee A, Kittur BH, Bargali SS, Bargali K, Nema S (2023) Forest biodiversity conservation and restoration: Policies, plan, and approaches. In: Banerjee A, Jhariya MK, Bargali SS, Palit D (eds) *Ecorestoration for Sustainability*, 1st edn. Wiley
- Ralph CJ, Sauer JR, Droege S (1995) Monitoring bird populations by point counts. *Gen Tech Rep PSW-GTR-149*. Albany, CA: U.S. Department of Agriculture, Forest Service, Pacific Southwest Research Station.
- Reed J, Van Vianen J, Deakin EL, Barlow J, Sunderland T (2016) Integrated landscape approaches to managing social and environmental issues in the tropics: learning from the past to guide the future. *Glob Chang Biol* 22:2540–2554. <https://doi.org/10.1111/gcb.13284>
- Republic of Rwanda (2012) Rwanda Vision 2020. [https://climatechange.gov.rw/fileadmin/user\\_upload/Documents/Report/RwandaVision2020.pdf](https://climatechange.gov.rw/fileadmin/user_upload/Documents/Report/RwandaVision2020.pdf) Accessed 12 June 2025
- Republic of Rwanda (2020) Gishwati-Mukura National Park named a Biosphere Reserve. [https://www.rema.gov.rw/info/details?tx\\_news\\_pi1%5Baction%5D=detail&tx\\_news\\_pi1%5Bcontroller%5D=News&tx\\_news\\_pi1%5Bnews%5D=94&cHash=c4137e09291abbc43f1844fe541d306a](https://www.rema.gov.rw/info/details?tx_news_pi1%5Baction%5D=detail&tx_news_pi1%5Bcontroller%5D=News&tx_news_pi1%5Bnews%5D=94&cHash=c4137e09291abbc43f1844fe541d306a) Accessed 11 June 2025
- Republic of Rwanda (2025) LAFREC. <https://www.rema.gov.rw/our-work/projects/lafrec> Accessed 12 June 2025
- Ricketts TH (2001) The matrix matters: effective isolation in fragmented landscapes. *Am Nat* 158:87–99. <https://doi.org/10.1086/320863>
- Rizinjirabake F, Mukamugema J, Ndagijimana JN, Murwanashyaka T, Mizero A (2024) Ethnobotany survey of medicinal plants in home gardens around Volcanoes National Park, Rwanda. *Rwanda J Eng, Sci, Technol Environ*. <https://doi.org/10.4314/rjeste.v6i1.5>
- Rooduijn B, Bongers F, Van Der Wal H (2018) Wild native trees in tropical homegardens of Southeast Mexico: fostered by fragmentation, mediated by management. *Agric Ecosyst Environ* 254:149–161. <https://doi.org/10.1016/j.agee.2017.10.015>
- Rudnick D, Ryan SJ, Beier P, Cushman SA, Dieffenbach F, Epps C, Gerber LR, Hartter JN, Jenness JS, Kintsch J, Merenlender AM, Perkl RM, Perziosi DV, Trombulack SC (2012) The role of landscape connectivity in planning and implementing conservation and restoration priorities. *Issues Ecol* 16:1–20
- Ruffell J, Clout MN, Didham RK (2017) The matrix matters, but how should we manage it? Estimating the amount of high-quality matrix required to maintain biodiversity in fragmented landscapes. *Ecography* 40:171–178. <https://doi.org/10.1111/ecog.02097>
- Ruticumugambi JA, Kaplin B, Blondeel H et al (2024) Diversity and composition of agroforestry species

- in two agro-ecological zones of Rwanda. *Agroforest Syst* 98:1421–1443. <https://doi.org/10.1007/s10457-024-01011-9>
- Sala OE, Stuart Chapin F III, Armesto JJ, Berlow E, Bloomfield J, Dirzo R, Huber-Sanwald E, Huenneke LF, Jackson RB, Kinzig A, Leemans R, Lodge DM, Mooney HA, Oesterheld M, Poff NL, Sykes MT, Walker BH, Walker M, Wall DH (2000) Global biodiversity scenarios for the year 2100. *Science* 287:1770–1774. <https://doi.org/10.1126/science.287.5459.1770>
- Santos PZF, Crouzeilles R, Sansevero JBB (2019) Can agroforestry systems enhance biodiversity and ecosystem service provision in agricultural landscapes? A meta-analysis for the Brazilian Atlantic Forest. *For Ecol Manag* 433:140–145. <https://doi.org/10.1016/j.foreco.2018.10.064>
- Sayer J, Chokkalingam U, Poulsen J (2004) The restoration of forest biodiversity and ecological values. *For Ecol Manag* 201:3–11. <https://doi.org/10.1016/j.foreco.2004.06.008>
- Seburanga JL (2013) Decline of indigenous crop diversity in colonial and postcolonial Rwanda. *Int J Biodivers*. <https://doi.org/10.1155/2013/401938>
- Sirko W, Kashubin S, Ritter M, Annkah A, Bouchareb YSE, Dauphin Y, Keyzers D, Neumann M, Cisse M, Quinn J (2021) Continental-scale building detection from high resolution satellite imagery (version 2). arXiv. <https://doi.org/10.48550/ARXIV.2107.12283>
- Soemarwoto O (1987) Homegardens: A traditional agroforestry system with a promising future. In *Agroforestry: A decade of development*, pp 157–170
- Stanturf JA, Kleine M, Mansourian S, Parrotta J, Madsen P, Kant P, Burns J, Bolte A (2019) Implementing forest landscape restoration under the Bonn Challenge: a systematic approach. *Ann For Sci*. <https://doi.org/10.1007/s13595-019-0833-z>
- Stevenson T, Fanshawe J (2002) *Birds of East Africa: Kenya, Tanzania, Uganda, Rwanda*. Princeton University Press, Burundi
- Suding K, Higgs E, Palmer M, Callicott JB, Anderson CB, Baker M, Gutrich JJ, Hondula KL, LaFevor MC, Larson BMH, Randall A, Ruhl JB, Schwartz KZS (2015) Committing to ecological restoration. *Science* 348:638–640. <https://doi.org/10.1126/science.aaa4216>
- Sunwar S, Thornström C-G, Subedi A, Bystrom M (2006) Home gardens in western Nepal: opportunities and challenges for on-farm management of agrobiodiversity. *Biodivers Conserv* 15:4211–4238. <https://doi.org/10.1007/s10531-005-3576-0>
- Taylor PD, Fahrig L, Henein K, Merriam G (1993) Connectivity is a vital element of landscape structure. *Oikos* 68:571–573. <https://doi.org/10.2307/3544927>
- Tesfay H, Negash M (2025) Ethiopia: Enhancing Landscape Connectivity Through Agroforests. In: Lapin K, Oettel J, Braun M, Konrad H (eds) *Ecological Connectivity of Forest Ecosystems*. Springer, Cham
- Tscharntke T, Klein AM, Kruess A, Steffan-Dewenter I, Thies C (2005) Landscape perspectives on agricultural intensification and biodiversity – ecosystem service management. *Ecol Lett* 8:857–874. <https://doi.org/10.1111/j.1461-0248.2005.00782.x>
- Tscharntke T, Milder JC, Schroth G et al (2015) Conserving biodiversity through certification of tropical agroforestry crops at local and landscape scales. *Conserv Lett* 8:14–23. <https://doi.org/10.1111/conl.12110>
- Vande weghe JP, Vande weghe GR (2010) *Birds in Rwanda*. Rwanda Development Board.
- Vargas-Cárdenas F, Navarro-Sigüenza AG, Ceccon E (2024) The contribution of traditional-homegardens to bird conservation in human-modified landscapes. *Agrofor Syst* 98:2605–2616. <https://doi.org/10.1007/s10457-024-01054-y>
- Wang C, Zhang W, Li X, Wu J (2022) A global meta-analysis of the impacts of tree plantations on biodiversity. *Glob Ecol Biogeogr* 31:576–587. <https://doi.org/10.1111/geb.13440>
- World Bank (2014) Rwanda landscape approach to forest restoration and conservation project. <https://documents1.worldbank.org/curated/en/704801657067823071/pdf/Rwanda-Landscape-Approach-to-Forest-Restoration-and-Conservation-LAFREC-Project.pdf> Accessed 12 June 2025
- World Bank (2021) Climate Change Knowledge Portal. <https://climateknowledgeportal.worldbank.org/country/rwanda> Accessed 12 June 2025
- Yinebeb M, Lulekal E, Bekele T, Lemessa D (2022) Homegardens plant species richness and their use types have positive associations across agricultural landscapes of Northwest Ethiopia. *Glob Ecol Conserv*. <https://doi.org/10.1016/j.gecco.2022.e02342>
- Zanaga D, Van De Kerchove R, Daems D et al (2022) ESA WorldCover 10 m 2021 v200 (Version v200). Zenodo. <https://doi.org/10.5281/ZENODO.7254221>

**Publisher's Note** Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.